

HPV is so comman ANY sexually active person is likely come to in contact with it.

Whether you're a dirty rooster or relatively chaste, you're still likely to come in contact with HPV. Take steps to help protect yourself from genital warts and HPV-related cancers with your free* HPV immunisation.

Talk to Student Health today or visit bestshot.co.nz to find out more.

Immunise against
HPV CANCERS/

* GARDASIL® 9 is a prescription medicine, for females aged 9–45 years and males 9–26 years of age. It is a nine-valent recombinant human papillomavirus (HPV) vaccine. GARDASIL® 9 is indicated for 9–45 year-old females and 9–26 year-old males, for prevention of cervical, vulvar, vaginal, and anal cancer, precancerous or dysplastic lesions, genital warts or lesions, and infection caused by the HPV types in the vaccine. Each 0.5 mL dose contains the following HPV L1 Proteins by type: 6 (30 µg), 11 (40 µg), 13 (20 µg), 33 (20 µg), 45 (20 µg), and 58 (20 µg), GARDASIL® 9 is right for you. Tell healthcare professional if you or your child have ever had an allergic reaction to any vaccine or to any of the listed ingredients for GARDASIL® 9, or if you or your child has a serious illness, blood disease, bleeding disorder, a high fever, or a weakened immune system (e.g. due to medicines, a genetic condition, or human immunodeficiency virus [HIV] infection). Common reactions are headache, fever, and injection-site pain, swelling, and redness. Fainting can occur. Allergic reactions are rare but serious. If you are worried by any side effects, see your doctor, pharmacist, or healthcare professional. Go to the nearest hospital if you experience wheezing, shortness of breath, or difficulty breathing; or pinkish, itchy swellings and rash, especially on the face. Immunisation does not replace the need for regular cervical cancer screening. Additional product information and Consumer Medicine Information is available from www.medsafe.govt.nz (October 2019), and from Seqirus (NZ) Ltd, Auckland, on 800 502 757. GARDASIL® 9 is funded for both males and females aged 9–26 years – normal charges will apply for other patients. Copyright © 2019 Merck Sharp & Dohme Corp., a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ, USA. All Rights Reserved.TAPS NA11653 NZ/GAR9/0120/0073. INSIGHT 9721A



Whatever you call the thing, make sure you keep HPV out of it.

Show your appreciation for your favourite little friend by helping keep him safe from HPV-related cancers and genital warts. HPV immunisation is free* from the Student Health Clinic, so get yourself down there right now or visit bestshot.co.nz to find out more.



* GARDASIL® 9 is a prescription medicine, for females aged 9–45 years and males 9–26 years of age. It is a nine-valent recombinant human papillomavirus (HPV) vaccine. GARDASIL® 9 is indicated for 9–45 year-old females and 9–26 year-old males, for prevention of cervical, vulvar, vaginal, and anal cancer, precancerous or dysplastic lesions, genital warts or lesions, and infection caused by the HPV types in the vaccine. Each 0.5 mL dose contains the following HPVL1 Proteins by type: 6 (30 µg), 11 (40 µg), 16 (60 µg), 18 (40 µg), 31 (20 µg), 31 (20 µg), 45 (20 µg), 45 (20 µg), 6ARDASIL® 9 has risks and benefits, and should be used strictly as directed. Ask your doctor if GARDASIL® 9 is right for you. Tell your healthcare professional if you or your child have ever had an allergic reaction to any vaccine or to any of the listed ingredients for GARDASIL® 9, or if you or your child have ever had an allergic reaction to any vaccine or to any of the listed ingredients for GARDASIL® 9, or if you or your child have ever had an allergic reaction to any vaccine or to any of the listed ingredients for GARDASIL® 9, or if you or your child have sever had an allergic reaction to any vaccine or to any of the listed ingredients for GARDASIL® 9, or if you or your child have sever had an allergic reaction to any vaccine or to any of the listed ingredients for GARDASIL® 9, or if you or your child have sever had an allergic reaction to any vaccine or to any of the listed ingredients for GARDASIL® 11 you or your child have sever had an allergic reaction to any vaccine or to any of the listed ingredients for GARDASIL® 11 you or your child have sever had nallergic reaction to any vaccine or to any of the listed ingredients for GARDASIL® 11 you or your child have sever had nallergic reaction to any vaccine or to any or the listed ingredients for the males and temales aged 9–26 years – normal charges will apply for other patients. Copyright © 2019 Merck Sharp & Dohme Corp., a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ, USA, A



BARRY THOMPSON BSC. HTV

Make sure a good education is the only thing you're picking up at Uni.

Learning the ways of the world is great, getting HPV while you're doing it is not. Help protect yourself from genital warts and HPV-related cancers with your free* HPV immunisation.

Talk to Student Health today or visit bestshot.co.nz to find out more.

Immunise against

HPV CANCERS/

*GARDASIL® 9 is a prescription medicine, for females aged 9–45 years and males 9–26 years of age. It is a nine-valent recombinant human papillomavirus (HPV) vaccine. GARDASIL® 9 is indicated for 9–45 year-old females and 9–26 year-old males, for prevention of cervical, vulvar, vaginal, and anal cancer, precancerous or dysplastic lesions, genital warts or lesions, and infection caused by the HPV types in the vaccine. Each 0.5 mL dose contains the following HPV L1 Proteins by type: 6 (30 µg), 11 (40 µg), 13 (20 µg), 33 (20 µg), 45 (20 µg), 52 (20 µg), and 58 (20 µg), GARDASIL® 9 has risks and benefits, and should be used strictly as directed. Ask your doctor if GARDASIL® 9 is right for you. Tell your healthcare professional if you or your child have ever had an allergic reaction to any vaccine or to any of the listed ingredients for GARDASIL® 9, or if you or your child has a serious illness, blood disease, bleeding disorder, a high fever, or weakened immune system (e.g. due to medicines, a genetic condition, or human immunodefficiency virus [HIV] infection). Common reactions are headache, fever, and injection-site pain, swelling, and redness. Fainting can occur. Altergic reactions are rare but serious. If you are worried by any side effects, see your doctor, pharmacist, or healthcare professional. Go to the nearest hospital if you experience wheezing, shortness of breath, or difficulty breathing; or pinkish, litchy swellings and rash, especially on the face. Immunisation does not replace the need for regular cervical cancer screening. Additional product information and Consumer Medicine Information is available from www.medsafe.govt.nz (October 2019), and from Sequins (NZ) Ltd, Auckland, on 800 502 757. GARDASIL® 918 (Indeed for both makes and females aged 9–26 years – normal charges will apply for other patients. Copyright © 2019 Merck Sharp & Dohme Corp., a subsidiary of Merck & Co., Inc., Michiebrose States of NA 1850 All Richel Research 1850 All Richel Research 1850 All Richel Research 1850 All Richel Researc